

NGC 1883: a neglected intermediate-age open cluster located in the outskirts of the Galactic disk[★]

Giovanni Carraro¹, Gustavo Baume^{1,2} and Sandro Villanova¹ [†]

¹*Dipartimento di Astronomia, Università di Padova, Vicolo dell'Osservatorio 2, I-35122 Padova, Italy*

²*Facultad de Ciencias Astronómicas y Geofísicas de la UNLP, IALP-CONICET, Paseo del Bosque s/n, La Plata, Argentina*

Submitted: March 2003

ABSTRACT

We report on *BVI* CCD photometry of a field centered in the region of the open cluster NGC 1883 down to $V = 21$. This cluster has never been studied insofar, and we provide for the first time estimates of its fundamental parameters, namely radial extent, age, distance and reddening. We find that the cluster has a radius of about 2.5 arcmin, and shows signatures of dynamical relaxation. NGC 1883 is located in the anti-center direction, and exhibits a reddening in the range $E(B - V) = 0.23 - 0.35$, depending on the metal abundance. It turns out to be of intermediate-age (1 billion years old), and quite distant for an open cluster. In fact it is located 4.8 kpc from the Sun, and more than 13 kpc from the Galactic center. This results makes NGC 1883 one of the most peripheral old open clusters, with important consequences for the trend of the metallicity with distance in the outer Galactic disk.

Key words: Open clusters and associations: general – open clusters and associations: individual: NGC 1883 - Hertzsprung-Russell (HR) diagram

1 INTRODUCTION

NGC 1883 (C 0522+465, OCL 417) is a northern open cluster, located toward the anti-center direction ($\alpha = 05^{\text{h}} 25^{\text{m}} .9$, $\delta = +46^{\circ} 29'$, $l = 163^{\circ} .08$, $b = +06^{\circ} .16$, J2000.). This cluster was never studied insofar. The only information we have come from Trumpler (1931) who suggests that this cluster is quite well detached from the field, shows a clear central concentration with a radius of about 3 arcmin, and is moderately rich (*I*2m Trumpler class). The same kind of information can be derived by consulting Collinder (1931) work. This paper presents the first photometric study of the cluster, and it is part of a series dedicated at improving the photometry of northern open clusters at Asiago Observatory (see Carraro et al. 2002, and references therein)

The plan of the paper is as follows. Sect. 2 illustrates the observation and reduction strategies. An analysis of the geometrical structure and star counts in the field of NGC 1883 are presented in Sect. 3, whereas a discussion of the Color-Magnitude Diagram (CMD) is performed in Sect. 4. Sect. 5

deals with the determination of cluster reddening, distance and age. Finally, Sect. 6 summarizes our findings.

2 OBSERVATIONS AND DATA REDUCTION

CCD *BVI* observations were carried out with the AFOSC camera at the 1.82 m Copernico telescope of Cima Ekar (Asiago, Italy), in the photometric night of November 8, 2002. AFOSC, with a pixel size of $0''.473$, samples a $8'.14 \times 8'.14$ field in a $1K \times 1K$ nitrogen-cooled thinned CCD.

Details of the observations are listed in Table 2 where the observed field is reported together with the exposure times, the typical seeing values and the air-masses. Fig. 1 shows the finding chart NGC 1883 region taken from the DSS-2[‡] archive. The data has been reduced with the IRAF[§] packages CCDRED, DAOPHOT, and PHOTCAL using the point spread function (PSF) method (Stetson 1987). The calibration equations obtained by observing Landolt (1992)

[★] Based on observations carried out at Mt Ekar, Asiago, Italy. All the photometry is available at WEBDA database: <http://obswww.unige.ch/webda/navigation.html>


[†] email: giovanni.carraro@unipd.it (GC); baume@pd.astro.it (GB); villanova@pd.astro.it (SV)

[‡] Second generation Digitized Sky Survey, <http://cadwww.dao.nrc.ca/cadcbin/getdss>

[§] IRAF is distributed by NOAO, which are operated by AURA under cooperative agreement with the NSF.

Table 1. Journal of observations of NGC 1883 and standard star fields together with calibration coefficients (November 8, 2002).

Field	Filter	Exposure time [sec.]		Seeing [μ]		Airmass
NGC 1883	B	600	60	5	2.1	1.015
	V	300	30	3	2.0	1.018
	I	300	30	3	2.0	1.020
PG 0231+051	B	300			2.4	1.324
	V	60			2.2	1.316
	I	90			2.2	1.315
PG 2213-006	B	150			2.3	1.291
	V	30			2.3	1.298
	I	30			2.5	1.295
Calibration coefficients	$b_1 = +1.602 \pm 0.004$					
	$b_2 = +0.038 \pm 0.006$					
	$b_3 = +0.29$					
	$v_{1bv} = +1.003 \pm 0.014$			$i_1 = +1.691 \pm 0.044$		
	$v_{2bv} = -0.016 \pm 0.018$			$i_2 = +0.057 \pm 0.043$		
	$v_3 = +0.16$			$i_3 = +0.08$		
	$v_{1vi} = +1.002 \pm 0.016$					
	$v_{2vi} = -0.013 \pm 0.016$					

**Figure 1.** A DSS image of a region around NGC 1883 covered by the present study. North is up, east on the left.

PG 0231+051 and PG 2213-006 fields observed along the night, are:



$$b = B + b_1 + b_2(B - V) + b_3X \quad (1)$$

$$v = V + v_{1bv} + v_{2bv}(B - V) + v_3X \quad (2)$$

$$v = V + v_{1vi} + v_{2vi}(V - I) + v_3X \quad (3)$$

$$i = I + i_1 + i_2(V - I) + i_3X \quad (4)$$

where BVI are standard magnitudes, bvi are the instrumental ones, X is the airmass and the derived coefficients are presented at the bottom of Table 2. As for V magnitudes, when B magnitude was available, we used expression (2) to

**Figure 2.** Photometric errors as a function of magnitude, for our NGC 1883 observations.**Figure 3.** Star counts in the field of NGC 1883 as a function of the radius.

compute them, elsewhere expression (3) was used. The standard stars in these fields provide a very good color coverage. For the extinction coefficients, we assumed the typical values for the Asiago Observatory (Desidera et al. 2002).

Finally, Fig. 2 presents the run of photometric errors as a function of magnitude. These errors take into account




Figure 4. CMDs of NGC 1883 as a function of the radius.

fitting errors from DAOPHOT and calibration errors, and have been computed following Patat & Carraro (2001). It can be noticed that stars brighter than $V \approx 21$ mag have photometric errors lower than 0.10 mag in magnitude and lower than 0.15 mag in colour. The final photometric data are available in electronic form at the WEBDA[¶] site.

3 STAR COUNTS AND CLUSTER SIZE

Lyngå (1987) and Dias et al. (2002) report a preliminary estimate of NGC 1883 radius, amounting to 2.5-3.0 arcmin. This is basically confirmed by the appearance of the cluster in the finding chart (Fig. 1), from which one can readily see that we were able to cover all the cluster region. We perform star counts by using our CCD data to obtain a new estimate of the cluster size. We derive the surface stellar density by performing star counts in concentric rings around stars #12

(the present numbering) selected as the approximate cluster center, and then dividing by their respective surfaces. The final density profile and the corresponding Poisson error bars are shown in Fig. 3, where we take into account all the measured stars brighter than $V \approx 19.5$ mag.

The surface density has a value of 10-12 stars/arcmin² within 2.5 arcmin, and a value of 4-6 stars/arcmin² outside this radius. From this sharp density variation we argue that the cluster population is dominant within 2.5 arcmin, whereas outside the general Galactic disk field dominates the star counts (see also Fig. 1).

The precise value of the cluster radius is however difficult to be obtained, and we consider 2.5 arcmin as a lower limit. Obviously, we expect that some cluster stars are located out of this radius, which we simply consider as the distance from the cluster center where the cluster dominates over the field.

4 THE COLOUR-MAGNITUDE DIAGRAMS

The CMDs of NGC 1883 are shown in Fig. 4. To facilitate the interpretation we consider the distribution of the stars

[¶] <http://obswww.unige.ch/webda/navigation.html>

in the CMD as a function of the distance from the cluster center. In the left panel of Fig. 4 we plot all the detected stars. Here the cluster exhibits a well defined Main Sequence (MS) extending from $V=20$ to $V=14.5$. Interestingly, there are a few stars in the red part of the CMD, which resemble a Red Giant Branch (RGB). The MS is significantly wide, and some hints for a substantial population of unresolved binary stars is visible - particularly in the upper MS - in the form of a parallel redder sequence. The Turn Off Point (TO) is roughly located at $V=16$, $(B-V)=0.60$. The group of stars above the TO (between $V=12.5$ and $V=13.5$) can be either blue stragglers belonging to the cluster or field stars. Finally the brightest star (see also Fig. 1), located southwest of the cluster center, and recorded as GSC 03358-01188 and TYC 3358 -1188 -1, is quite probably an interloper. Much better information can be obtained by looking at the middle and right panels in Fig. 4. The middle panel contains only the stars located inside the estimated cluster radius (2.5 arcmin), whereas the right panel contains all the stars located outside the cluster radius. The following considerations can be done:

- The MS and the TO region in the middle panel are much better defined;
- almost all the probable RGB stars are inside the inner region, which implies by the way that the cluster underwent some dynamical relaxation;
- the stars above the TO are probably field stars, since they lie all out of the cluster radius (see right panel).

In particular the fine shape of the TO deserves some attention. In fact the shape of the TO is that one typical of intermediate-age open clusters, with a blue and red hook clearly visible notwithstanding the field stars contamination. This is a clear indication of an age in the range 1-1.5 Gyr, depending on the precise metal content of the cluster (Carraro & Chiosi 1994, Carraro et al. 1999).

5 CLUSTER FUNDAMENTAL PARAMETERS

In this section we try to provide some estimates of the cluster basic parameters. To achieve this result, we make use of the comparison between the stars distribution in the CMDs and a set of theoretical isochrones from the Padova group (Girardi et al. 2000). We already have an indication of the cluster age, but we do not know anything about the reddening, the distance and the metallicity. The results of the fitting are shown in Fig. 5 and 6.

In details, in Fig. 5 we present the CMDs for the stars within 2.5 arcmin from the cluster center, with over-imposed an isochrone of 1.2 billion years for a solar ($Z=0.019$) metallicity. The fit is quite good in the V vs $(B-V)$ plane (left panel), but the shape of the MS is not very well reproduced in the V vs $(V-I)$ plane (right panel). Moreover the precise location of the RGB is not well matched. We achieved this results by shifting the isochrone with $E(B-V) = 0.23$, $E(V-I) = 0.30$, and $(m-M) = 14.00$ and by adopting $R_V = 3.1$.

We tried to improve the fit by lowering the metal abundance, on the base of the fact that a cluster located toward the anti-center might have a lower than solar metal abundance (Friel 1995). The result is shown in Fig. 6. Here the isochrone has




Figure 5. **Left panel:** NGC 1883 data in the V vs. $B-V$ diagram, as compared to Girardi et al. (2000) isochrone of age 1.2×10^9 yr (solid line), for the metallicity $Z = 0.019$. A distance modulus of $(m-M)_0 = 13.25$ mag, and a colour excess of $E(B-V) = 0.23$ mag, have been derived. **Right panel:** NGC 1883 data in the V vs. $V-I$ diagram, as compared to Girardi et al. (2000) isochrone of age 1.2×10^9 yr (solid line), for the metallicity $Z = 0.019$. A distance modulus of $(m-M)_0 = 13.15$ mag, and a colour excess of $E(V-I) = 0.30$ mag, have been derived.

an age of 1.0 billion years for the $Z=0.008$ metal content. The global fit is quite good, and has been obtained by shifting the isochrone with $E(B-V) = 0.35$, $E(V-I) = 0.50$, and $(m-M) = 14.50$ ($R_V = 3.1$).

In conclusion, a lower than solar metal abundance seems to be favored. If this is the case, NGC 1883 turns out to be located 4.8 kpc from the Sun toward the anti-center direction. This implies a distance from the Galactic center somewhat larger than 13 kpc, by assuming $R_\odot = 8.5$ kpc, and a height above the Galactic plane of about 500 pc. According to Friel (1995, Table 1), NGC 1883 turns out to be one of the few intermediate-age or old open clusters located in the outskirts of the Galactic disk. In fact we know only other 3 clusters (Tombaugh 2, Berkeley 20 and Berkeley 29) located more distant than NGC 1883. Therefore NGC 1883 might play an important role in defining the precise shape of the radial abundance gradient in the external regions of the Galactic disk, which is one of the fundamental constraints of models of chemical evolution (Chiappini et al. 2001). Moreover NGC 1883 lies significantly above the Galactic plane. The position, together with its compactness, are probably the main reasons for which the cluster survived until the present time.

6 CONCLUSIONS

We have presented the first CCD BVI photometric study of the open cluster NGC 1883. The CMDs we derive allow us




Figure 6. **Left panel:** NGC 1883 data in the V vs. $B - V$ diagram, as compared to Girardi et al. (2000) isochrone of age 1.0×10^9 yr (solid line), for the metallicity $Z = 0.008$. A distance modulus of $(m - M)_0 = 13.40$ mag, and a colour excess of $E(B - V) = 0.35$ mag, have been derived. **Right panel:** NGC 1883 data in the V vs. $V - I$ diagram, as compared to Girardi et al. (2000) isochrone of age 1.0×10^9 yr (solid line), for the metallicity $Z = 0.008$. A distance modulus of $(m - M)_0 = 13.20$ mag, and a colour excess of $E(V - I) = 0.50$ mag, have been derived.

to infer estimates of the cluster basic parameters. In detail, we find that:

- the age of NGC 1883 is around 1.0 Gyr;
- the reddening E_{B-V} turns out to be 0.23 ± 0.10 by adopting a metal abundance $Z=0.019$, or 0.35 ± 0.10 mag if one assumes $Z=0.008$;
- we place the cluster at about 4.8 kpc from the Sun toward the anti-center direction;
- this way NGC 1883 turns out to be one of the most distant intermediate-age open cluster.

Future work should concentrate to obtain an estimate of the cluster metal abundance through spectra of the RGB stars. The knowledge of the cluster metallicity, which we could not constrain very well, is of paramount importance to better probe the trend of the metallicity in the outer parts of the Galactic disk (Friel & Janes 1993).

ACKNOWLEDGEMENTS

We acknowledge the kind night assistance by Asiago Observatory technical staff and in particular dr. Stefano Ciroi who secured part of the observations discussed in this work. This study has been financed by the Italian Ministry of University, Scientific Research and Technology (MURST) and the Italian Space Agency (ASI), and made use of Simbad and WEBDA databases.

REFERENCES

- Carraro G., Chiosi C., 1994, A&A 287, 761
 Carraro G., Barbon R., Boschetti C.S., 2002, MNRAS 336, 259
 Carraro G., Girardi L., Chiosi C., 1999, MNRAS 309, 430
 Carraro G., Ng Y.K., Portinari L., 1998, MNRAS 296, 1045
 Chiappini C., Matteucci F., Romano D., 2001, ApJ 554, 1044
 Collinder P., 1931, Ld. An., 2
 Desidera S., Fantinel D., Giro E., Afosc User Manual, 2002
 Dias W.S., Alessi B.S., Moitinho A., Lepine J.R.D., 2002, A&A 389, 871
 Friel E.D., 1995, ARA&A 33, 381
 Friel E.D., Janes K.A., 1993, A&A 267, 75
 Friel E.D., Janes K.A., Tavaréz M., et al., 2002, AJ 124, 2693
 Girardi L., Bressan A., Bertelli G., Chiosi C., 2000, A&AS 141, 371
 Landolt A.U., 1992, AJ 104, 340
 Lyngå G., 1987, The Open Star Clusters Catalogue, 5th edition
 Patat F., Carraro G., 2001, MNRAS 325, 1591
 Trumpler R.J., 1930, Lick Observ. Bull. 14, 154